

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/51984564>

2-(4-Fluorophenyl)-2-oxoethyl 3-(trifluoromethyl)benzoate

Article in *Acta Crystallographica Section E Structure Reports Online* · November 2011

DOI: 10.1107/S160053681103947X · Source: PubMed

CITATIONS

7

READS

44

5 authors, including:



Suhana Arshad

Universiti Sains Malaysia

336 PUBLICATIONS 358 CITATIONS

[SEE PROFILE](#)



Bhadrachar Garudachar

Kuwait Institute for Scientific Research

152 PUBLICATIONS 358 CITATIONS

[SEE PROFILE](#)



Arun M Islor

National Institute of Technology Karnataka

548 PUBLICATIONS 4,208 CITATIONS

[SEE PROFILE](#)



Shivananda K N

Technion - Israel Institute of Technology

75 PUBLICATIONS 493 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Desalination [View project](#)



Medicinal organic chemistry [View project](#)

2-(4-Fluorophenyl)-2-oxoethyl 3-(trifluoromethyl)benzoate

Hoong-Kun Fun,^{a*}‡ Suhana Arshad,^a B. Garudachari,^b
A. M. Isloor^b and Kammasandra N Shivananda^c

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, ^bMedicinal Chemistry Division, Department of Chemistry, National Institute of Technology-Karnataka, Surathkal, Mangalore 575 025, India, and ^cSchulich Faculty of Chemistry, Technion Israel Institute of Technology, Haifa 32000, Israel
Correspondence e-mail: hkfun@usm.my


Received 23 September 2011; accepted 26 September 2011

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; disorder in main residue; R factor = 0.057; wR factor = 0.182; data-to-parameter ratio = 21.8.

In the title compound, $\text{C}_{16}\text{H}_{10}\text{F}_4\text{O}_3$, the fluoroform group is disordered over two orientations with an occupancy ratio of 0.834 (4):0.166 (4). The dihedral angle between the two aromatic rings is $20.34(9)^\circ$. In the crystal, $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into layers lying parallel to the bc plane.

Related literature

For background to the chemistry of phenacyl benzoate derivatives, see: Huang *et al.* (1996); Gandhi *et al.* (1995); Ruzicka *et al.* (2002); Litera *et al.* (2006); Sheehan & Umezawa (1973). For bond-length data, see: Allen *et al.* (1987). For a related structure, see: Fun *et al.* (2011).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{10}\text{F}_4\text{O}_3$
 $M_r = 326.24$
Monoclinic, $P2_1/c$
 $a = 14.7694(19)\text{ \AA}$
 $b = 12.1602(16)\text{ \AA}$
 $c = 8.0929(10)\text{ \AA}$
 $\beta = 95.886(2)^\circ$
 $V = 1445.8(3)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.14\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.38 \times 0.25 \times 0.07\text{ mm}$

Data collection

Bruker SMART APEXII DUO
CCD diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2009)
 $T_{\min} = 0.950$, $T_{\max} = 0.990$
18790 measured reflections
4815 independent reflections
2841 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.182$
 $S = 1.04$
4815 reflections
221 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.34\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.35\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C5—H5A \cdots O3 ⁱ	0.93	2.50	3.263 (2)	139
C8—H8A \cdots O1 ⁱⁱ	0.97	2.55	3.502 (2)	169

Symmetry codes: (i) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HKF and SA thank Universiti Sains Malaysia for the Research University Grants (No. 1001/PFIZIK/811160). SA thanks the Malaysian Government and USM for the Academic Staff Training Scheme (ASTS) award. AMI thanks Professor Sandeep Sanchethi, Director, National Institute of Technology-Karnataka, India for providing research facilities and the Board for Research in Nuclear Sciences, Department of Atomic Energy, Government of India for the Young Scientist award.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6418).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bruker (2009). *SADABS*, *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Fun, H.-K., Arshad, S., Garudachari, B., Isloor, A. M. & Satyanarayana, M. N. (2011). *Acta Cryst. E67*, o1528.
Gandhi, S. S., Bell, K. L. & Gibson, M. S. (1995). *Tetrahedron*, **51**, 13301–13308.
Huang, W., Pian, J., Chen, B., Pei, W. & Ye, X. (1996). *Tetrahedron*, **52**, 10131–10136.
Litera, J. K., Loya, A. D. & Klan, P. (2006). *J. Org. Chem.* **71**, 713–723.
Ruzicka, R., Zabadal, M. & Klan, P. (2002). *Synth. Commun.* **32**, 2581–2590.
Sheehan, J. C. & Umezawa, K. (1973). *J. Org. Chem.* **38**, 3771–3773.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

‡ Thomson Reuters ResearcherID: A-3561-2009.

supplementary materials

Acta Cryst. (2011). E67, o2836 [doi:10.1107/S160053681103947X]

2-(4-Fluorophenyl)-2-oxoethyl 3-(trifluoromethyl)benzoate

H.-K. Fun, S. Arshad, B. Garudachari, A. M. Isloor and K. N. Shivananda

Comment

In organic chemistry, phenacyl benzoate is a derivative of an acid formed by reaction between acid and phenacyl bromide. They find applications in the field of synthetic chemistry (Huang *et al.*, 1996; Gandhi *et al.*, 1995) such as synthesis of oxazoles, imidazoles and benzoxazepines. They are also useful for photo-removable protecting groups for carboxylic acids in organic synthesis and biochemistry (Ruzicka *et al.*, 2002; Litera *et al.*, 2006; Sheehan & Umezaw, 1973). Keeping this in view, the title compound was synthesized to study its crystal structure.

In the molecular structure (Fig. 1), the fluoro form group is disordered over two orientations with an occupancy ratio of 0.834 (4):0.166 (4). The two phenyl rings (C1–C6 & C10–C15) make a dihedral angle of 20.34 (9) $^{\circ}$. Bond lengths (Allen *et al.*, 1987) and angles are within normal range and are comparable to the related structures (Fun *et al.*, 2011).

The crystal packing is shown in Fig. 2. Intermolecular C5—H5A \cdots O3 and C8—H8A \cdots O1 hydrogen bonds (Table 1) linked the molecules into layers parallel to *bc* plane.

Experimental

A mixture of 3-(trifluoromethyl)benzoic acid (1.0 g, 0.0052 mol), potassium carbonate (0.78 g, 0.0057 mol) and 2-bromo-1-(4-fluorophenyl)ethanone (1.12 g, 0.0052 mol) in dimethylformamide (10 ml) was stirred at room temperature for 2 h. On cooling, colourless needle-shaped crystals of 2-(4-fluorophenyl)-2-oxoethyl 3-(trifluoromethyl)benzoate begin to separate out. These were collected by filtration and recrystallized from ethanol to yield colourles blocks. Yield: 1.6 g, 93.5%. *M.p.*: 369–370 K.

Refinement

The fluoro form group is disordered over two orientations, with a final refined occupancy ratio of 0.834 (4):0.166 (4). All H atoms were positioned geometrically [C–H = 0.93 or 0.97 Å] and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids. Both disordered components are shown.

supplementary materials




Fig. 2. The crystal packing of the title compound. The dashed lines represent the hydrogen bonds. Only the major disordered components are shown.

2-(4-Fluorophenyl)-2-oxoethyl 3-(trifluoromethyl)benzoate

Crystal data

C ₁₆ H ₁₀ F ₄ O ₃	<i>F</i> (000) = 664
<i>M_r</i> = 326.24	<i>D_x</i> = 1.499 Mg m ⁻³
Monoclinic, <i>P</i> 2 ₁ /c	Mo <i>K</i> α radiation, λ = 0.71073 Å
Hall symbol: -P 2ybc	Cell parameters from 4165 reflections
<i>a</i> = 14.7694 (19) Å	θ = 2.2–27.2°
<i>b</i> = 12.1602 (16) Å	μ = 0.14 mm ⁻¹
<i>c</i> = 8.0929 (10) Å	<i>T</i> = 296 K
β = 95.886 (2)°	Block, colourless
<i>V</i> = 1445.8 (3) Å ³	0.38 × 0.25 × 0.07 mm
<i>Z</i> = 4	

Data collection

Bruker SMART APEXII DUO CCD diffractometer	4815 independent reflections
Radiation source: fine-focus sealed tube graphite	2841 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.028$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2009)	$\theta_{\text{max}} = 31.6^\circ$, $\theta_{\text{min}} = 1.4^\circ$
$T_{\text{min}} = 0.950$, $T_{\text{max}} = 0.990$	$h = -21 \rightarrow 21$
18790 measured reflections	$k = -17 \rightarrow 17$
	$l = -11 \rightarrow 11$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.057$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.182$	H-atom parameters constrained
$S = 1.04$	$w = 1/[\sigma^2(F_o^2) + (0.0745P)^2 + 0.3289P]$
4815 reflections	where $P = (F_o^2 + 2F_c^2)/3$
221 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
	$\Delta\rho_{\text{max}} = 0.34 \text{ e \AA}^{-3}$

0 restraints

 $\Delta\rho_{\min} = -0.35 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
F1	0.41559 (16)	0.37665 (19)	-0.5044 (2)	0.1144 (9)	0.834 (4)
F2	0.44559 (19)	0.25120 (16)	-0.3326 (4)	0.1179 (11)	0.834 (4)
F3	0.53719 (16)	0.3931 (3)	-0.3405 (5)	0.1359 (12)	0.834 (4)
F1X	0.5096 (11)	0.4001 (13)	-0.4146 (18)	0.110 (5)*	0.166 (4)
F2X	0.4251 (10)	0.2913 (17)	-0.413 (2)	0.137 (6)*	0.166 (4)
F3X	0.5187 (7)	0.2898 (8)	-0.2308 (12)	0.096 (3)*	0.166 (4)
F4	-0.27124 (10)	0.35096 (15)	0.5827 (2)	0.1118 (6)	
O1	0.18686 (10)	0.27074 (10)	-0.0004 (2)	0.0756 (4)	
O2	0.15270 (8)	0.43444 (10)	0.10304 (15)	0.0578 (3)	
O3	0.02656 (9)	0.56904 (9)	0.18644 (16)	0.0623 (3)	
C1	-0.11779 (12)	0.51838 (14)	0.3751 (2)	0.0522 (4)	
H1A	-0.1071	0.5922	0.3543	0.063*	
C2	-0.18848 (13)	0.48938 (17)	0.4648 (2)	0.0628 (5)	
H2A	-0.2257	0.5427	0.5050	0.075*	
C3	-0.20243 (14)	0.38007 (19)	0.4931 (3)	0.0707 (5)	
C4	-0.14948 (14)	0.29857 (17)	0.4363 (3)	0.0720 (5)	
H4A	-0.1609	0.2250	0.4577	0.086*	
C5	-0.07894 (13)	0.32825 (14)	0.3467 (2)	0.0577 (4)	
H5A	-0.0423	0.2741	0.3071	0.069*	
C6	-0.06199 (11)	0.43833 (12)	0.31499 (18)	0.0447 (3)	
C7	0.01478 (11)	0.47363 (12)	0.22220 (19)	0.0458 (3)	
C8	0.07752 (12)	0.38408 (13)	0.1730 (2)	0.0523 (4)	
H8A	0.0994	0.3408	0.2696	0.063*	
H8B	0.0451	0.3356	0.0921	0.063*	
C9	0.20286 (12)	0.36718 (13)	0.0189 (2)	0.0520 (4)	
C10	0.27900 (11)	0.42548 (13)	-0.0491 (2)	0.0502 (4)	
C11	0.30026 (13)	0.53430 (15)	-0.0112 (3)	0.0658 (5)	
H11A	0.2676	0.5727	0.0624	0.079*	
C12	0.36985 (16)	0.58564 (18)	-0.0826 (3)	0.0818 (7)	
H12A	0.3833	0.6589	-0.0581	0.098*	
C13	0.41962 (15)	0.52905 (18)	-0.1900 (3)	0.0753 (6)	

supplementary materials

H13A	0.4665	0.5639	-0.2379	0.090*
C14	0.39943 (13)	0.42070 (16)	-0.2259 (2)	0.0604 (4)
C15	0.32944 (12)	0.36853 (14)	-0.1566 (2)	0.0550 (4)
H15A	0.3161	0.2953	-0.1820	0.066*
C16	0.45169 (18)	0.3601 (2)	-0.3439 (4)	0.0866 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.153 (2)	0.1219 (18)	0.0750 (12)	0.0103 (13)	0.0431 (12)	-0.0150 (11)
F2	0.162 (2)	0.0723 (12)	0.1333 (19)	0.0433 (13)	0.0827 (18)	0.0078 (12)
F3	0.0656 (13)	0.194 (3)	0.156 (3)	-0.0032 (14)	0.0486 (15)	-0.064 (2)
F4	0.0863 (10)	0.1217 (13)	0.1375 (13)	-0.0144 (9)	0.0607 (9)	0.0101 (10)
O1	0.0871 (10)	0.0428 (7)	0.1035 (11)	-0.0067 (6)	0.0413 (8)	-0.0073 (7)
O2	0.0601 (7)	0.0439 (6)	0.0729 (8)	-0.0040 (5)	0.0234 (6)	-0.0052 (5)
O3	0.0732 (8)	0.0360 (6)	0.0797 (8)	-0.0019 (5)	0.0170 (6)	0.0077 (5)
C1	0.0571 (9)	0.0416 (8)	0.0576 (9)	0.0070 (7)	0.0040 (7)	-0.0007 (7)
C2	0.0567 (10)	0.0668 (12)	0.0655 (10)	0.0109 (9)	0.0099 (8)	-0.0044 (9)
C3	0.0556 (11)	0.0800 (14)	0.0791 (13)	-0.0082 (10)	0.0185 (9)	0.0035 (10)
C4	0.0748 (13)	0.0512 (10)	0.0933 (14)	-0.0120 (9)	0.0253 (11)	0.0056 (10)
C5	0.0640 (10)	0.0377 (8)	0.0735 (11)	-0.0022 (7)	0.0171 (8)	-0.0004 (7)
C6	0.0494 (8)	0.0350 (7)	0.0493 (8)	0.0001 (6)	0.0033 (6)	0.0004 (6)
C7	0.0542 (9)	0.0338 (7)	0.0491 (8)	-0.0008 (6)	0.0042 (6)	0.0008 (6)
C8	0.0580 (9)	0.0404 (8)	0.0606 (9)	-0.0023 (7)	0.0164 (7)	0.0015 (7)
C9	0.0572 (9)	0.0425 (8)	0.0574 (9)	0.0022 (7)	0.0117 (7)	0.0004 (7)
C10	0.0503 (9)	0.0429 (8)	0.0581 (9)	0.0028 (6)	0.0090 (7)	0.0009 (7)
C11	0.0630 (11)	0.0483 (9)	0.0894 (13)	-0.0005 (8)	0.0236 (10)	-0.0124 (9)
C12	0.0803 (14)	0.0515 (11)	0.1196 (18)	-0.0159 (10)	0.0387 (13)	-0.0176 (11)
C13	0.0658 (12)	0.0634 (12)	0.1012 (16)	-0.0088 (10)	0.0307 (11)	-0.0018 (11)
C14	0.0562 (10)	0.0586 (10)	0.0685 (11)	0.0035 (8)	0.0172 (8)	-0.0001 (8)
C15	0.0596 (10)	0.0446 (8)	0.0623 (10)	0.0038 (7)	0.0129 (8)	-0.0012 (7)
C16	0.0847 (16)	0.0780 (16)	0.1050 (19)	-0.0056 (13)	0.0472 (14)	-0.0146 (14)

Geometric parameters (\AA , $^\circ$)

F1—C16	1.368 (4)	C5—C6	1.391 (2)
F2—C16	1.331 (3)	C5—H5A	0.9300
F3—C16	1.323 (3)	C6—C7	1.486 (2)
F1X—C16	1.182 (16)	C7—C8	1.510 (2)
F2X—C16	1.059 (17)	C8—H8A	0.9700
F3X—C16	1.537 (10)	C8—H8B	0.9700
F4—C3	1.355 (2)	C9—C10	1.482 (2)
O1—C9	1.203 (2)	C10—C11	1.387 (2)
O2—C9	1.336 (2)	C10—C15	1.387 (2)
O2—C8	1.4342 (19)	C11—C12	1.379 (3)
O3—C7	1.2126 (18)	C11—H11A	0.9300
C1—C2	1.377 (3)	C12—C13	1.379 (3)
C1—C6	1.395 (2)	C12—H12A	0.9300
C1—H1A	0.9300	C13—C14	1.375 (3)

C2—C3	1.368 (3)	C13—H13A	0.9300
C2—H2A	0.9300	C14—C15	1.380 (2)
C3—C4	1.371 (3)	C14—C16	1.484 (3)
C4—C5	1.377 (3)	C15—H15A	0.9300
C4—H4A	0.9300		
C9—O2—C8	115.59 (13)	C12—C11—C10	120.07 (17)
C2—C1—C6	120.80 (16)	C12—C11—H11A	120.0
C2—C1—H1A	119.6	C10—C11—H11A	120.0
C6—C1—H1A	119.6	C13—C12—C11	120.44 (18)
C3—C2—C1	118.20 (17)	C13—C12—H12A	119.8
C3—C2—H2A	120.9	C11—C12—H12A	119.8
C1—C2—H2A	120.9	C14—C13—C12	119.56 (19)
F4—C3—C2	118.5 (2)	C14—C13—H13A	120.2
F4—C3—C4	118.41 (19)	C12—C13—H13A	120.2
C2—C3—C4	123.04 (18)	C13—C14—C15	120.65 (18)
C3—C4—C5	118.40 (18)	C13—C14—C16	119.73 (19)
C3—C4—H4A	120.8	C15—C14—C16	119.59 (19)
C5—C4—H4A	120.8	C14—C15—C10	119.88 (16)
C4—C5—C6	120.66 (17)	C14—C15—H15A	120.1
C4—C5—H5A	119.7	C10—C15—H15A	120.1
C6—C5—H5A	119.7	F2X—C16—F1X	108.5 (12)
C5—C6—C1	118.91 (15)	F2X—C16—F3	123.5 (9)
C5—C6—C7	122.18 (14)	F1X—C16—F2	120.0 (8)
C1—C6—C7	118.90 (14)	F3—C16—F2	111.8 (3)
O3—C7—C6	122.18 (14)	F2X—C16—F1	61.8 (11)
O3—C7—C8	121.35 (14)	F1X—C16—F1	73.2 (7)
C6—C7—C8	116.47 (12)	F3—C16—F1	104.7 (3)
O2—C8—C7	108.49 (12)	F2—C16—F1	100.9 (3)
O2—C8—H8A	110.0	F2X—C16—C14	122.9 (9)
C7—C8—H8A	110.0	F1X—C16—C14	123.9 (8)
O2—C8—H8B	110.0	F3—C16—C14	113.3 (2)
C7—C8—H8B	110.0	F2—C16—C14	114.0 (2)
H8A—C8—H8B	108.4	F1—C16—C14	111.1 (2)
O1—C9—O2	123.44 (16)	F2X—C16—F3X	93.2 (11)
O1—C9—C10	124.35 (16)	F1X—C16—F3X	93.6 (8)
O2—C9—C10	112.20 (14)	F3—C16—F3X	66.5 (4)
C11—C10—C15	119.39 (16)	F2—C16—F3X	56.6 (4)
C11—C10—C9	122.52 (15)	F1—C16—F3X	144.3 (4)
C15—C10—C9	118.08 (15)	C14—C16—F3X	103.9 (4)
C6—C1—C2—C3	0.0 (3)	C15—C10—C11—C12	1.2 (3)
C1—C2—C3—F4	179.32 (18)	C9—C10—C11—C12	-177.7 (2)
C1—C2—C3—C4	-0.1 (3)	C10—C11—C12—C13	-0.9 (4)
F4—C3—C4—C5	-179.30 (19)	C11—C12—C13—C14	0.0 (4)
C2—C3—C4—C5	0.1 (3)	C12—C13—C14—C15	0.6 (3)
C3—C4—C5—C6	0.0 (3)	C12—C13—C14—C16	178.9 (2)
C4—C5—C6—C1	-0.1 (3)	C13—C14—C15—C10	-0.3 (3)
C4—C5—C6—C7	178.56 (17)	C16—C14—C15—C10	-178.6 (2)
C2—C1—C6—C5	0.2 (2)	C11—C10—C15—C14	-0.6 (3)

supplementary materials


C2—C1—C6—C7	−178.58 (15)	C9—C10—C15—C14	178.38 (16)
C5—C6—C7—O3	175.79 (16)	C13—C14—C16—F2X	−154.8 (13)
C1—C6—C7—O3	−5.5 (2)	C15—C14—C16—F2X	23.5 (14)
C5—C6—C7—C8	−3.7 (2)	C13—C14—C16—F1X	−2.1 (9)
C1—C6—C7—C8	174.96 (14)	C15—C14—C16—F1X	176.3 (9)
C9—O2—C8—C7	−165.10 (14)	C13—C14—C16—F3	32.0 (4)
O3—C7—C8—O2	7.2 (2)	C15—C14—C16—F3	−149.7 (3)
C6—C7—C8—O2	−173.24 (13)	C13—C14—C16—F2	161.3 (3)
C8—O2—C9—O1	0.9 (3)	C15—C14—C16—F2	−20.3 (4)
C8—O2—C9—C10	−179.97 (14)	C13—C14—C16—F1	−85.5 (3)
O1—C9—C10—C11	−173.9 (2)	C15—C14—C16—F1	92.9 (3)
O2—C9—C10—C11	7.0 (2)	C13—C14—C16—F3X	102.0 (5)
O1—C9—C10—C15	7.2 (3)	C15—C14—C16—F3X	−79.6 (5)
O2—C9—C10—C15	−171.99 (15)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C5—H5A…O3 ⁱ	0.93	2.50	3.263 (2)	139
C8—H8A…O1 ⁱⁱ	0.97	2.55	3.502 (2)	169

Symmetry codes: (i) $-x, y-1/2, -z+1/2$; (ii) $x, -y+1/2, z+1/2$.

Fig. 1



supplementary materials

Fig. 2

